



King's Research Portal

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Chiossi, S., & Salamon, S. (2002). The intrinsic torsion of $SU(3)$ and G_2 structures. In *Differential Geometry, Valencia*.

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

THE INTRINSIC TORSION OF $SU(3)$ AND G_2 STRUCTURES

SIMON CHIOSSI

*Dipartimento di Matematica, Università di Genova, Via Dodecaneso 35,
I-16146 Genova*

SIMON SALAMON

*Dipartimento di Matematica, Politecnico di Torino, Corso Duca degli Abruzzi 24,
I-10129 Torino*

To Antonio Naveira on the occasion of his 60th birthday

We analyse the relationship between the components of the intrinsic torsion of an $SU(3)$ -structure on a 6-manifold and a G_2 -structure on a 7-manifold. Various examples illustrate the type of $SU(3)$ -structure that can arise as a reduction of a metric with holonomy G_2 .

Introduction

Let G be a subgroup of $SO(n)$. A G -structure on a smooth manifold M of dimension n induces a Riemannian metric g on M . The failure of the holonomy group of the Levi-Civita connection of g to reduce to G is measured by the so-called intrinsic torsion τ . It is known^{1,2} that the latter is a tensor which takes values at each point in $T^* \otimes \mathfrak{g}^\perp$ where T^* is the cotangent space and \mathfrak{g}^\perp is the orthogonal complement of \mathfrak{g} in $\mathfrak{so}(n) \cong \bigwedge^2 T^*$.

This note is concerned with the cases

- (1) $SU(3) \subset SO(6)$,
- (2) $G_2 \subset SO(7)$.

The respective tensors τ_1 and τ_2 belong to spaces of dimension 42 and 49. The fact that $SU(3)$ is a maximal subgroup of G_2 gives a direct relationship between the two structures. Indeed, the sets of reductions (1) and (2) are both parametrized by the projective space

$$\mathbb{RP}^7 = \frac{SO(6)}{SU(3)} = \frac{SO(7)}{G_2}. \quad (1)$$

The fact that this space itself admits homogeneous G_2 -structures has applications to the study of families of G_2 -structures. Moreover, the fibration $\mathbb{RP}^7 \rightarrow \mathbb{CP}^3$ is indicative of the way in which G_2 -structures can in general be built from almost-Hermitian structures on a 6-manifold.

We begin by describing the tensor τ_1 determined by an $SU(3)$ -structure on a 6-manifold M , thereby refining the theory for $U(3)$. An additional summand in the $SU(3)$ case can be used to construct a new conformally invariant torsion tensor. It is well known that a holonomy reduction to $SU(3)$ is characterized by the existence of a symplectic form together with a closed form of ‘type $(3, 0)$ ’, and it follows that all the components of τ_1 can be calculated in terms of *exterior* derivatives of the forms defining the reduction. The special relevance of 3-forms in describing 6-dimensional structures is already documented,³ and this paper presents some additional applications.

In the general set-up, the $SU(3)$ reduction leads to a splitting of the Nijenhuis tensor in two equal parts, which give rise to different components of the tensor τ_2 on a 7-manifold \mathbb{M} whose structure reduces from G_2 to $SU(3)$. This aspect of the theory is reminiscent of self-duality in four dimensions, and the G_2 examples analysed in subsequent sections can by analogy be divided into those of self-dual and anti-self-dual type. The distinction arises from whether the 7-manifold is foliated by leaves of dimension 1 or 6.

Our first descriptions of τ_1, τ_2 are ‘static’ in the sense that they relate to a fixed G -structure and are purely algebraic. We subsequently examine how the components of τ_1 determine those of τ_2 in various situations in which the geometry of the 6- and 7-manifolds are interrelated, with the inclusion $SU(3) \subset G_2$ varying from point to point. The evolution equations discussed by Hitchin⁴ are interpreted using the notion of a half-flat $SU(3)$ structure. We provide additional examples of incomplete metrics with holonomy G_2 of the type discovered by Gibbons et al⁵ that suggest that half-flat structures occur naturally on 6-dimensional nilmanifolds.

The final section undertakes an investigation of certain cases in which the G_2 -manifold \mathbb{M} is a circle bundle over a 6-manifold M endowed with an appropriate structure. We provide an explicit description of τ_2 as a function of τ_1 and a curvature 2-form, and consider the case of the canonical circle bundle over a Kähler 3-fold. When the holonomy of \mathbb{M} reduces to G_2 , the quotient $M = \mathbb{M}/S^1$ is a symplectic manifold with a type of generalized Calabi-Yau geometry that we describe briefly in terms of τ_1 . Examples of such quotients of the known complete metrics with G_2 holonomy incorporate interesting global features,^{6,7} and it is hoped that the techniques of this paper will aid a fuller investigation of this situation.

1 Static $SU(3)$ structures

Let M be a 6-manifold with a $U(3)$ -structure. Thus M is equipped with a Riemannian metric g , an orthogonal almost-complex structure J and an associated 2-form ω . The exterior forms on M may be decomposed into types relative to J , and we adopt the following notation⁸ at each point:

$$\begin{aligned} T^* &= [\Lambda^{1,0}], \\ \wedge^2 T^* &= [\Lambda^{2,0}] \oplus [\Lambda^{1,1}] \cong [\Lambda^{2,0}] \oplus [\Lambda_0^{1,1}] \oplus \mathbb{R}, \\ \wedge^3 T^* &= [\Lambda^{3,0}] \oplus [\Lambda^{2,1}] \cong [\Lambda^{3,0}] \oplus [\Lambda_0^{2,1}] \oplus [\Lambda^{1,0}], \\ \wedge^4 T^* &= [\Lambda^{3,1}] \oplus [\Lambda^{2,2}] \cong [\Lambda^{2,0}] \oplus [\Lambda_0^{1,1}] \oplus \mathbb{R}. \end{aligned} \quad (2)$$

The induced metric distinguishes the circle B consisting of elements of unit norm in the 2-dimensional space $[\Lambda^{3,0}]$. An $SU(3)$ -structure is determined by the choice of a real 3-form ψ_+ lying in B at each point, or equivalently a section of the associated S^1 -bundle \mathbb{B} . The associated $(3,0)$ -form is

$$\Psi = 2(\psi_+)^{3,0} = \psi_+ + i\psi_- \quad (3)$$

with $\psi_- = J\psi_+$. We may then write

$$\mathbb{B} = \{a\psi_+ + b\psi_- : a^2 + b^2 = 1\}, \quad (4)$$

and this description remains valid locally even if a global reduction from $U(3)$ to $SU(3)$ does not exist.

To be more explicit, we may choose a local orthonormal basis (e^1, \dots, e^6) of T^* such that

$$\Psi = (e^1 + ie^2) \wedge (e^3 + ie^4) \wedge (e^5 + ie^6).$$

Consequently

$$\begin{aligned} \omega &= e^{12} + e^{34} + e^{56}, \\ \psi_+ &= e^{135} - e^{146} - e^{236} - e^{245}, \\ \psi_- &= e^{136} + e^{145} + e^{235} - e^{246}, \end{aligned} \quad (5)$$

where e^{135} stands for $e^1 \wedge e^3 \wedge e^5$ etc. These forms are subject to the compatibility relations

$$\begin{aligned} \omega \wedge \psi_{\pm} &= 0, \\ \psi_+ \wedge \psi_- &= \frac{2}{3}\omega^3. \end{aligned} \quad (6)$$

Table 1. $SU(3)$ torsion

component	$\dim_{\mathbb{R}}$	$U(3)$ -module	$SU(3)$ -module	
\mathcal{W}_1	2	$[\![\Lambda^{3,0}]\!]$	\mathbb{R}	\mathbb{R}
\mathcal{W}_2	16	$[\![V]\!]$	$\mathfrak{su}(3)$	$\mathfrak{su}(3)$
\mathcal{W}_3	12	$[\![\Lambda_0^{2,1}]\!]$	$[\![S^{2,0}]\!]$	
\mathcal{W}_4	6	T	T	
\mathcal{W}_5	6	T	T	

The intrinsic torsion of the $U(3)$ -structure can be identified with ∇J or $\nabla\omega$ and belongs to the space

$$T^* \otimes \mathfrak{u}(3)^\perp = \mathcal{W}_1 \oplus \mathcal{W}_2 \oplus \mathcal{W}_3 \oplus \mathcal{W}_4,$$

whose four components were first described by Gray–Hervella.⁹ The intrinsic torsion τ_1 of the $SU(3)$ -structure lies in the enlarged space

$$T^* \otimes \mathfrak{su}(3)^\perp \cong T^* \otimes ([\![\Lambda^{2,0}]\!] \oplus \mathbb{R}) \cong T \otimes (T \oplus \mathbb{R}),$$

given that now $\Lambda^{2,0} \cong \Lambda^{0,1}$. Thus

$$\tau_1 \in \mathcal{W}_1 \oplus \mathcal{W}_2 \oplus \mathcal{W}_3 \oplus \mathcal{W}_4 \oplus \mathcal{W}_5,$$

with $\mathcal{W}_5 \cong T$. Properties of the various components are indicated by Table 1.

We denote the component of τ_1 in \mathcal{W}_i in a formal way by W_i , though we shall need to supply more precise definitions shortly. It is well known that the components of ∇J in $\mathcal{W}_1 \oplus \mathcal{W}_3 \oplus \mathcal{W}_4$ are determined by $d\omega$, and those in $\mathcal{W}_1 \oplus \mathcal{W}_2$ by the Nijenhuis tensor. The choice of basis (ψ_+, ψ_-) of $[\![\Lambda^{3,0}]\!]$ at each point provides an isomorphism

$$\mathcal{W}_1 \oplus \mathcal{W}_2 = [\![\Lambda^{2,0} \otimes \Lambda^{1,0}]\!] = [\![\Lambda^{3,0} \otimes \Lambda^{1,1}]\!] \cong \mathbb{R}^2 \otimes \mathfrak{u}(3).$$

The rank of $W_1 + W_2$ (or of W_2 on its own) in this tensor product is equal to one of 0, 1, 2.

Remark. If M is an almost-Hermitian manifold for which the bundle \mathbb{B} is trivial but not actually trivialized, the basis (ψ_+, ψ_-) is defined up to an overall constant action by S^1 , and the rank of $W_1 + W_2$ is a global invariant. This situation occurs naturally on 6-dimensional Lie groups of the type considered below.

On a complex manifold, $d\psi_+$ belongs to $\Lambda^{3,1} \oplus \Lambda^{1,3}$ at each point. It follows that the component of $d\psi_+$ in $\Lambda^{2,2}$ is determined by the Nijenhuis tensor, and therefore by $W_1 \oplus W_2$. We may define the two scalar components $W_1^\pm \in \mathbb{R}$ of W_1 by

$$\begin{aligned} d\psi_+ \wedge \omega &= \psi_+ \wedge d\omega = W_1^+ \omega^3, \\ d\psi_- \wedge \omega &= \psi_- \wedge d\omega = W_1^- \omega^3, \end{aligned}$$

where $\omega^3 = \omega \wedge \omega \wedge \omega$. Similarly, $W_2 = W_2^+ + W_2^-$ in which

$$\begin{aligned} (d\psi_+)^{2,2} &= W_1^+ \omega^2 + W_2^+ \wedge \omega, \\ (d\psi_-)^{2,2} &= W_1^- \omega^2 + W_2^- \wedge \omega, \end{aligned}$$

so that $W_2^\pm \in [\Lambda_0^{1,1}]$ are effective $(1,1)$ -forms.

Given that

$$d\psi_+ - id\psi_- = d\bar{\Psi} \in \Lambda^{1,3} \oplus \Lambda^{2,2},$$

the remaining components of $d\psi_+, d\psi_-$ are related by

$$(d\psi_+)^{3,1} = i(d\psi_-)^{3,1}. \quad (7)$$

It is now clear that the W_5 -component of τ_1 arises from (7). In summary, we have

1.1 Theorem The five components of τ_1 are determined by $d\omega, d\psi_+, d\psi_-$, in the following manner:

$$\begin{aligned} W_1 &\longleftrightarrow (d\omega)^{3,0} \\ W_2 &\longleftrightarrow ((d\psi_+)_0^{1,1}, (d\psi_-)_0^{1,1}) \\ W_3 &\longleftrightarrow (d\omega)_0^{2,1} \\ W_4 &\longleftrightarrow \omega \wedge d\omega \\ W_5 &\longleftrightarrow (d\psi_\pm)^{3,1} \end{aligned}$$

(refer to (2)).

It is significant that W_4 and W_5 arise from isotypic summands of the space $T^* \otimes \mathfrak{su}(3)^\perp$. Before moving on to seven dimensions, it is convenient to give a more precise definition of these components too in order that they may be compared directly. We shall do this by means of the contraction

$$\lrcorner : \bigwedge^k T^* \otimes \bigwedge^n T^* \rightarrow \bigwedge^{n-k} T^*$$

that exploits the underlying Riemannian metric, with the convention that $e^{12} \lrcorner e^{12345} = e^{345}$ etc.

1.2 Definition The components of τ_1 in $\mathcal{W}_4, \mathcal{W}_5$ are given by

$$\begin{aligned} W_4 &= \frac{1}{2}\omega \lrcorner d\omega, \\ W_5 &= \frac{1}{2}\psi_+ \lrcorner d\psi_+. \end{aligned}$$

The coefficient of one half is added with the following examples in mind. Given ω as in (5), suppose that $d\omega = \omega \wedge e^1$ and $d\psi_+ = \psi_+ \wedge e^1$. Then

$$\begin{aligned} W_4 &= \frac{1}{2}\omega \lrcorner (e^{134} + e^{156}) = e^1, \\ W_5 &= \frac{1}{2}\psi_+ \lrcorner (e^{1236} + e^{1245}) = e^1. \end{aligned} \tag{8}$$

Now suppose that

$$(d\psi_+)^{3,1} = \Psi \wedge \bar{\sigma} = (\psi_+ + i\psi_-) \wedge \bar{\sigma}.$$

Then $W_5 = 2(\bar{\sigma} + \sigma)$, whereas $\frac{1}{2}\psi_+ \lrcorner d\psi_- = 2i(\sigma - \bar{\sigma})$ by (7). It follows that

$$\psi_+ \lrcorner d\psi_- = J(\psi_+ \lrcorner d\psi_+), \tag{9}$$

a useful re-interpretation of (7).

Each of the components W_1, W_2, W_3 is at worst re-scaled under a conformal change of metric

$$g \mapsto e^{2f}g. \tag{10}$$

This is a consequence of the fact that none of the corresponding representations in Table 1 is isomorphic to the cotangent space T^* containing the 1-form df . The reduction to $SU(3)$ permits one to define an additional conformally invariant component:

1.3 Proposition The tensor $3W_4 + 2W_5$ is unchanged by (10).

Proof. The transformation (10) multiplies 1-forms by e^f . Hence the exterior derivatives of ω, ψ_+ transform as

$$\begin{aligned} d\omega &\mapsto d(e^{2f}\omega) = e^{2f}d\omega + 2e^{2f}df \wedge \omega, \\ d\psi_+ &\mapsto d(e^{3f}\psi_+) = e^{3f}d\psi_+ + 3e^{3f}df \wedge \psi_+. \end{aligned}$$

Retaining \lrcorner exclusively for the contraction relative to the original metric,

$$\begin{aligned} W_4 &\mapsto W_4 + \omega \lrcorner (df \wedge \omega), \\ W_5 &\mapsto W_5 + \frac{3}{2}\psi_+ \lrcorner (df \wedge \psi_+). \end{aligned}$$

The final terms may be evaluated by using (8) with $df = e^1$, with the result that they cancel out in the sum $3W_4 + 2W_5$. \square

2 Static G_2 structures

We denote by \mathbb{T} the space \mathbb{R}^7 , regarded as the standard representation of the exceptional Lie group G_2 . The latter acts transitively on the sphere S^6 in \mathbb{T} , and the stabilizer of a point of S^6 is conjugate to a fixed subgroup $SU(3)$ of G_2 . The inclusion $SU(3) \subset G_2$ is therefore characterized by the orthogonal decomposition

$$\mathbb{T} = T \oplus \mathbb{R}. \quad (11)$$

We choose an orthonormal basis (e^i) of \mathbb{T}^* such that $\alpha = e^7$ annihilates T at each point.

A G_2 -structure on a 7-manifold \mathbb{M} is characterized by a ‘positive generic 3-form’ φ . Adopting a canonical form compatible with (5), we set

$$\begin{aligned} \varphi &= \omega \wedge \alpha + \psi_+ \\ &= e^{127} + e^{347} + e^{567} + e^{135} - e^{146} - e^{236} - e^{245}. \end{aligned} \quad (12)$$

The basis (e^i) is orthonormal for the metric determined by the inclusion $G_2 \subset SO(7)$, and allows us to consider

$$\begin{aligned} *\varphi &= \psi_- \wedge \alpha + \frac{1}{2}\omega^2 \\ &= e^{1367} + e^{1457} + e^{2357} - e^{2467} + e^{3456} + e^{1256} + e^{1234}. \end{aligned} \quad (13)$$

The structure of a general G_2 -manifold will not reduce to $SU(3)$, and these descriptions are only valid pointwise or locally.

The intrinsic torsion space

$$\mathbb{T}^* \otimes \mathfrak{g}_2^\perp = \mathcal{X}_1 \oplus \mathcal{X}_2 \oplus \mathcal{X}_3 \oplus \mathcal{X}_4$$

has four components of respective dimensions 1, 14, 27, 7, first described by Fernandez–Gray.¹⁰ Various constructions^{1,6,11,12} of metrics with holonomy equal to G_2 are based on the significant fact that the holonomy reduction is characterized by the simultaneous closure of φ and $*\varphi$.

Given that \mathbb{T} represents the tangent space of \mathbb{M} , the G_2 counterpart of (2) consists of the decompositions

$$\begin{aligned} \mathbb{T}^* &\cong \mathbb{T} \\ \bigwedge^2 \mathbb{T}^* &\cong \mathfrak{g}_2 \oplus \mathbb{T} \\ \bigwedge^3 \mathbb{T}^* &\cong \mathbb{R} \oplus \mathbb{T} \oplus S_0^2 \mathbb{T}. \end{aligned} \quad (14)$$

It follows that $\mathfrak{g}_2^\perp \cong \mathbb{T}$, and the spaces $\mathcal{X}_1, \mathcal{X}_2, \mathcal{X}_3, \mathcal{X}_4$ are isomorphic to $\mathbb{R}, \mathfrak{g}_2, S_0^2 \mathbb{T}, \mathbb{T}$ respectively. The components of $\nabla \varphi$ can be recovered from those

Table 2. G_2 torsion

component	dim	G_2 -module	$SU(3)$ -module		
\mathcal{X}_1	1	\mathbb{R}	\mathbb{R}		
\mathcal{X}_2	14	\mathfrak{g}_2	T		$\mathfrak{su}(3)$
\mathcal{X}_3	27	$S_0^2\mathbb{T}$	$\llbracket S^{2,0} \rrbracket$	\mathbb{R}	T $\mathfrak{su}(3)$
\mathcal{X}_4	7	\mathbb{T}	\mathbb{R}		T

of $d\varphi$ and $d*\varphi$ as follows:

$$\begin{aligned}
X_1 + X_3 + X_4 &\longleftrightarrow d\varphi, \\
X_1 &\longleftrightarrow d\varphi \wedge \varphi, \\
X_2 + X_4 &\longleftrightarrow d*\varphi, \\
X_4 &\longleftrightarrow (*d*\varphi) \wedge (*\varphi).
\end{aligned} \tag{15}$$

The G_2 -structure is called *calibrated* (respectively *cocalibrated*) if $d\varphi = 0$ (respectively $d*\varphi = 0$), and examples with all possible combinations of torsion components are known.^{13,14,15}

Whilst \mathcal{X}_4 is isomorphic to (11), there are analogous decompositions of the other spaces in (14), and therefore of \mathcal{X}_2 and \mathcal{X}_3 , under the action of $SU(3)$. These are described in Table 2. In particular, $S_0^2\mathbb{T}$ also contains a 1-dimensional $SU(3)$ -module, and it follows that

$$\begin{aligned}
&\alpha \\
&\omega \\
&\psi_+, \quad \omega \wedge \alpha, \quad \psi_- \\
&\psi_- \wedge \alpha, \quad \omega^2, \quad \psi_+ \wedge \alpha \\
&\omega^2 \wedge \alpha \\
&\omega^3
\end{aligned} \tag{16}$$

is a list of the exterior forms on \mathbb{M} fixed by $SU(3)$. With reference to the summands in (14), we may assert that

2.1 Lemma

$$\left. \begin{array}{l} \psi_- \in \mathbb{T} \\ 3\psi_+ - 4\omega \wedge \alpha \in S_0^2\mathbb{T} \end{array} \right\} \subset \bigwedge^3 \mathbb{T}^*, \quad \left. \begin{array}{l} \psi_+ \wedge \alpha \in \mathbb{T} \\ 3\psi_- \wedge \alpha - 2\omega^2 \in S_0^2\mathbb{T} \end{array} \right\} \subset \bigwedge^4 \mathbb{T}^*.$$

Proof. Given (6), $\gamma = 3\psi_+ - 4\omega \wedge \alpha$ satisfies $\gamma \wedge (*\varphi) = 0$ and is therefore orthogonal to φ . Since $-\psi_-$ is the contraction of $*\varphi$ with the tangent vector dual to α , it lies in the submodule \mathbb{T} of $\bigwedge^3 \mathbb{T}^*$. Since γ is also orthogonal to ψ_- , it lies in $S_0^2 \mathbb{T}$. The invariant 4-forms are obtained by observing that $*\psi_- = \psi_+ \wedge \alpha$ and $*(\omega \wedge \alpha) = \frac{1}{2}\omega^2$. \square

Remark. The existence of various $SU(3)$ -invariant elements of $\bigwedge^3 \mathbb{T}$ gives rise to a choice of induced G_2 -structures in the passage from 6 to 7 dimensions. For example,

$$3\psi_+ - 4\omega \wedge \alpha = 3 \left[\omega \wedge \left(-\frac{4}{3}\alpha\right) + \psi_+ \right]$$

determines a G_2 structure with reversed orientation on \mathbb{T} and different scalings relative to (11). Other choices of coefficients will have the effect of modifying combinations in Theorem 3.1 below.

The three components of $\mathbb{T}^* \otimes \mathfrak{g}_2^\perp$ isomorphic to T can be detected from corresponding components of $d\varphi$ and $d*\varphi$. It is useful to record the following list for diagnostic purposes.

2.2 Lemma

$$\begin{aligned} \zeta &= e^{1347} + e^{1567} - e^{1236} - e^{1245} \in T \subset \mathbb{T} \subset \bigwedge^4 \mathbb{T}^*, \\ \eta &= e^{1347} + e^{1567} + e^{1236} + e^{1245} \in T \subset S_0^2 \mathbb{T} \subset \bigwedge^4 \mathbb{T}^*, \\ \xi &= e^{13456} + e^{12357} - e^{12467} \in T \subset \mathbb{T} \subset \bigwedge^5 \mathbb{T}^*, \\ \vartheta &= 2e^{13456} - e^{12357} + e^{12467} \in T \subset \mathfrak{g}_2 \subset \bigwedge^5 \mathbb{T}^*. \end{aligned}$$

Proof. Each of these forms represents the element of T dual to e^1 in an appropriate guise. For example, $\zeta = e^1 \wedge \varphi$ and η is a linear combination of ζ and $\psi_+ \wedge e^1 = e^{1236} + e^{1245}$ orthogonal to ζ . We may define ξ as $e^1 \wedge (*\varphi) = *(e^1 \lrcorner \varphi)$. Then ϑ is a linear combination of ξ and $e^1 \wedge \omega^2 = 2e^{13456}$ orthogonal to ξ . \square

3 Product manifolds

We now suppose that \mathbb{M} is a 7-manifold with an $SU(3)$ -structure, so that the differential forms $\alpha, \omega, \psi_+, \psi_-$ of respective degrees 1, 2, 3, 3 and constant norm are all defined globally. In this and the following sections, we shall investigate properties of the G_2 -structure defined with the convention of (12). In general, one may write $d\alpha = \alpha \wedge \beta + \gamma$, where β, γ are forms with values

in the subspace T^* at each point. For example, the equation $\gamma = 0$ is the integrability condition for the 6-dimensional distribution now determined by (11). We shall consider various special cases, the simplest of which is that in which \mathbb{M} is the Riemannian product of M with an interval or circle, so that $\nabla\alpha$ (and so $d\alpha$) vanishes.

In the product situation, we choose to write $\alpha = e^7 = dt$, so that

$$\begin{aligned} d\varphi &= d\omega \wedge dt + d\psi_+, \\ d * \varphi &= d\psi_- \wedge dt + \omega \wedge d\omega. \end{aligned} \tag{17}$$

Let

$$\begin{aligned} \tau_1 &= (W_1^+ + W_1^-) + (W_2^+ + W_2^-) + W_3 + W_4 + W_5, \\ \tau_2 &= X_1 + X_2 + X_3 + X_4 \end{aligned}$$

denote the respective intrinsic torsion tensors, as defined in the previous sections. Since $\nabla\varphi$ can be computed in terms of $\nabla\omega$ and $\nabla\psi_+$, the tensor τ_2 is determined by τ_1 . Any $SU(3)$ -invariant component of τ_2 must be a linear combination of W_1^+ and W_1^- , and any component isomorphic to T a linear combination of W_4 and W_5 . The precise statement is

3.1 Theorem The four components of τ_2 are determined by the seven components of τ_1 as follows.

$$\begin{aligned} X_1 &\longleftrightarrow W_1^+ \\ X_2 &\longleftrightarrow (W_2^-, 2W_4 + W_5) \\ X_3 &\longleftrightarrow (W_1^+, W_2^+, W_3, W_4 + W_5) \\ X_4 &\longleftrightarrow (W_4 - W_5, W_1^-). \end{aligned}$$

Proof. The component X_1 arises from

$$\varphi \wedge d\varphi = \psi_+ \wedge d\omega \wedge dt + \omega \wedge dt \wedge d\psi_+ = 2W_1^+ \omega^3 \wedge \alpha.$$

Similarly, X_4 is determined by the contraction

$$\varphi \lrcorner d\varphi = -\omega \lrcorner d\omega + \psi_+ \lrcorner d\psi_+ + (\psi_+ \lrcorner d\omega)dt.$$

The first two terms of the right-hand side are $-2W_4$ and $2W_5$ by Definition 1.2, and the last term corresponds to $d\omega \wedge \psi_-$ (given that $\psi_+ \wedge \psi_+ = 0$) and so W_1^- . This justifies the description of X_4 .

The hypotheses $d\omega = \omega \wedge e^1$ and $d\psi_- = \psi_- \wedge e^1$ are compatible with the constraint $W_4 = W_5$. In this case

$$d * \varphi = -e^{12357} + e^{12467} + 2e^{13456} = \vartheta,$$

confirming that $X_4 = 0$. In order to obtain ξ instead of ϑ , we need to take $d\omega = \frac{1}{2}\omega \wedge e^1$ and $d\psi_- = -\psi_- \wedge e^1$, which corresponds to $2W_4 + W_5 = 0$.

The association of W_2^- with X_2 and W_2^+, W_3 with X_3 follows immediately from (15). The hypotheses $d\omega = \omega \wedge e^1$ and $d\psi_+ = -\psi_+ \wedge e^1$ are compatible with the constraint $W_4 + W_5 = 0$. This implies that

$$d\varphi = e^{1347} + e^{1567} - e^{1236} - e^{1245} = \zeta,$$

whence the T -component of X_3 is proportional to $W_4 + W_5$. \square

Remark. The difference $49 - 42 = 7$ of the dimensions of the spaces containing τ_1 and τ_2 is accounted for by the repetition of W_1^+ and a linear combination of W_4, W_5 in the above list. This redundancy is eliminated in the more complicated situations described in subsequent sections. The lack of repetition between components of X_1, X_2 is consistent with the result¹⁴ that a connected G_2 manifold with $\tau_2 \in \mathcal{X}_1 \oplus \mathcal{X}_2$ has at least one of X_1, X_2 zero.

3.2 Corollary Suppose that M has an $SU(3)$ -structure. The G_2 -structure defined on $M \times \mathbb{R}$ by (12) is cocalibrated if and only if $\tau_1 \in \mathcal{W}_2^-$.

Examples. 1. An almost-Hermitian 6-manifold is called nearly-Kähler¹⁶ if ∇J belongs to the space \mathcal{W}_1 . Assuming that $\nabla J \neq 0$, the structure reduces to $SU(3)$ and we may suppose that $\tau_1 \in \mathcal{W}_1^-$ with ψ_+ proportional to $d\omega$. The product $M \times S^1$ then has a G_2 -structure with $\tau_2 \in \mathcal{X}_4$. Alternatively we may swap the roles of ψ_+, ψ_- to obtain $\tau_2 \in \mathcal{X}_1 \oplus \mathcal{X}_3$.

2. A known example¹³ of a calibrated nilmanifold can be interpreted as follows. Let $\mathfrak{g} = \mathbb{R} \oplus \mathfrak{h}$ be a 6-dimensional Lie algebra with structure determined by

$$de^i = \begin{cases} 0, & i = 1, 2, 4, 5 \\ e^{25}, & i = 3, \\ -e^{24}, & i = 6. \end{cases}$$

The definitions (5) furnish an associated nilmanifold $M = \Gamma \backslash G$ with an $SU(3)$ -structure for which

$$d\omega = 0, \quad d\psi_+ = 0, \quad d\psi_- = e^{1234} - e^{1256},$$

whence $\tau_1 \in \mathcal{W}_2^-$. It follows that $M \times S^1$ has both a calibrated G_2 structure and (swapping ψ_+, ψ_-) a cocalibrated one with $\tau_2 \in \mathcal{X}_3$. The same Lie algebra \mathfrak{g} was incidentally used¹⁷ in the construction of a closed non-parallel 4-form with stabilizer $Sp(2)Sp(1)$.

4 Dynamic G_2 structures

Let M be a fixed 6-manifold. Suppose that (ω, ψ_+, ψ_-) is an $SU(3)$ structure that depends on a real parameter t lying in some interval $I \subseteq \mathbb{R}$, so that one may regard $\mathbb{M} = M \times I$ as a warped product fibring over I .

To avoid confusion, we denote exterior differentiation on M by \hat{d} in this section. Adopting a unit 1-form dt on I allows us to write

$$\begin{aligned} d\varphi &= (\hat{d}\omega - \frac{\partial\psi_+}{\partial t}) \wedge dt + \hat{d}\psi_+, \\ d*\varphi &= (\hat{d}\psi_- + \omega \wedge \frac{\partial\omega}{\partial t}) \wedge dt + \omega \wedge \hat{d}\omega. \end{aligned}$$

This motivates

4.1 Definition An almost Hermitian 6-manifold is *half-flat* if it possesses a reduction to $SU(3)$ for which $d\psi_+ = 0$ and $\omega \wedge d\omega = 0$.

Half-flatness is therefore characterized by the closure of ψ_+ and ω^2 . It amounts to requiring that $W_1 + W_2$ has rank one and that both W_4, W_5 vanish. This eliminates $1 + 8 + 6 + 6 = 21$ of the total 42 dimensions of τ_1 , which is constrained to lie in $\mathcal{W}_1^- \oplus \mathcal{W}_2^- \oplus \mathcal{W}_3$.

If we now suppose that the G_2 -structure on \mathbb{M} has holonomy group contained in G_2 , we may conclude that M is half-flat for all t , and that the forms evolve according to the equations

$$\begin{cases} \hat{d}\omega = \frac{\partial\psi_+}{\partial t}, \\ \hat{d}\psi_- = -\omega \wedge \frac{\partial\omega}{\partial t}. \end{cases} \quad (18)$$

Conversely, suppose we are given a half-flat $SU(3)$ -structure at time $t = t_0$. The equations (18) may be regarded as a system constraining a closed 3-form ψ_+ and a closed 4-form ω^2 . For, if (as in our situation) the stabilizer of ψ_+ is $SL(3, \mathbb{C})$ then ψ_+ determines ψ_- via (3). In this way, Hitchin⁴ proved that the compatibility equations (6) are conserved in time, and this leads to

4.2 Theorem Let M be an almost Hermitian 6-manifold which is half-flat. Then there exists a metric with holonomy contained in G_2 on $M \times I$ for some interval I .

A key example of this construction is the following. Given (M, \hat{g}) , consider the conical metric $g = t^2\hat{g} + dt^2$ on $M \times \mathbb{R}^+$. Consistent with this, we set

$$\omega = t^2\hat{\omega}, \quad \psi_+ = t^3\hat{\psi}_+, \quad \psi_- = t^3\hat{\psi}_-,$$

where circumflex indicates a form independent of t . Then (18) becomes

$$d\hat{\omega} = 3\hat{\psi}_+, \quad d\hat{\psi}_- = -2\hat{\omega}^2.$$

These equations determine the nearly-Kähler class for which $\tau_1 \in \mathcal{W}_1^-$.

Further examples. 1. We now construct metrics with holonomy G_2 associated to each of the three nilmanifolds with $b_1 = 4$ whose $U(3)$ structures were classified by Abbena et al.¹⁸ Starting with the Iwasawa manifold M , we modify the usual basis of 1-forms in order that

$$de^i = \begin{cases} 0, & i = 1, 2, 3, 4, \\ -e^{14} - e^{23}, & i = 5, \\ -e^{13} - e^{42}, & i = 6. \end{cases} \quad (19)$$

The metric

$$g = t^2 \sum_{i=1}^4 e^i \otimes e^i + t^{-2} \sum_{i=5}^6 e^i \otimes e^i + t^4 dt^2$$

is compatible with the natural fibration $M \rightarrow T^4$, and the forms

$$\begin{aligned} \omega &= t^2(e^{12} + e^{34}) + t^{-2}e^{56}, \\ \psi_+ + i\psi_- &= t(e^1 + ie^2) \wedge (e^3 + ie^4) \wedge (e^5 + ie^6) \end{aligned}$$

determine a reduction to $SU(3)$ for which

$$\hat{d}\omega = t^{-3}\psi_+, \quad \hat{d}\psi_- = -4te^{1234},$$

whence $\tau_1 \in \mathcal{W}_1^- \oplus \mathcal{W}_2^-$. Indeed, g is a metric with holonomy G_2 on $M \times \mathbb{R}^+$.

2. The previous example was discovered by Gibbons et al⁵ who exhibit it as arising from the standard complete metric with holonomy G_2 by ‘contracting’ the isometry group $SO(5)$. A more complicated 2-step example is associated to the Lie algebra with

$$de^i = \begin{cases} 0, & i = 1, 2, 3, 4, \\ -e^{14} - e^{23}, & i = 5, \\ e^{24}, & i = 6. \end{cases}$$

Consider an orthonormal basis of 1-forms

$$te^1, \quad t^2e^2, \quad te^3, \quad t^2e^4, \quad t^{-2}e^5, \quad t^{-1}e^6, \quad 2t^4dt,$$

and the $SU(3)$ -structure for which

$$\begin{aligned}\omega &= t^3(e^{12} + e^{34}) + t^{-3}e^{56}, \\ \psi_+ + i\psi_- &= (e^1 + ite^2) \wedge (e^3 + ite^4) \wedge (e^5 + ite^6),\end{aligned}$$

by analogy to (5). This yields a G_2 -structure with closed forms

$$\begin{aligned}\varphi &= 2t^7(e^{12} + e^{34}) \wedge dt + 2te^{56} \wedge dt + e^{135} - t^2(e^{146} + e^{236} + e^{245}), \\ *\varphi &= -2t^7e^{246} \wedge dt + 2t^5(e^{145} + e^{136} + e^{235}) \wedge dt + e^{1256} + e^{3456} + t^6e^{1234}.\end{aligned}$$

3. The nilpotent 3-step Lie algebra for which

$$de^i = \begin{cases} 0, & i = 1, 2, 4, 5 \\ e^{25}, & i = 3, \\ e^{14} - e^{23}, & i = 6 \end{cases}$$

gives rise to an example satisfying the hypotheses of Theorem 4.2. In fact, the forms (5) satisfy

$$\omega \wedge d\omega = 0, \quad d\psi_+ = 0, \quad d\psi_- = -e^{1256}.$$

An explicit determination of the metric requires an analysis of spaces of invariant exact forms.⁴

5 Fibred G_2 manifolds

Let (M, \hat{g}) be a Riemannian 6-manifold. In this final section, we consider the 7-dimensional total space of a circle fibration $\pi : \mathbb{M} \rightarrow M$, endowed with a metric of the form

$$g = \alpha \otimes \alpha + \pi^*\hat{g}, \quad (20)$$

with $d\alpha = \pi^*\rho$ for some 2-form ρ on M .

To begin with, suppose that M has an $SU(3)$ -structure. The G_2 -structure on \mathbb{M} determined by (12) satisfies the following enhanced version of (17), in which we omit the pullback operator π^* :

$$\begin{aligned}d\varphi &= d\omega \wedge \alpha + d\psi_+ + \omega \wedge \rho. \\ d*\varphi &= d\psi_- \wedge \alpha + \omega \wedge d\omega - \psi_- \wedge \rho.\end{aligned} \quad (21)$$

Indicating the components of the 2-form ρ by

$$\begin{aligned}\bigwedge^2 T^* &= \langle \omega \rangle \oplus [\Lambda_0^{1,1}] \oplus [\Lambda^{2,0}] \\ \rho &= \rho_0\omega + \rho_1 + \rho_2,\end{aligned}$$

we obtain the following generalization of Theorem 3.1.

5.1 Theorem The four components of τ_2 are given by

$$\begin{aligned} X_1 &\longleftrightarrow W_1^+ + \rho_0 \\ X_2 &\longleftrightarrow (W_2^-, 2W_4 + W_5 - 2\rho_2) \\ X_3 &\longleftrightarrow (3W_1^+ - 4\rho_0, W_2^+ + \rho_1, W_3, W_4 + W_5 + \rho_2) \\ X_4 &\longleftrightarrow (W_4 - W_5 - \rho_2, W_1^-). \end{aligned}$$

Proof. Whilst X_1 corresponds to $\varphi \wedge d\varphi$, Lemma 2.1 tells us that the 1-dimensional component in X_3 arises from $(3\omega \wedge \alpha - 4\psi_+) \wedge d\varphi$. The various coefficients of ρ_2 can be deduced from the observations: (i) $d\varphi = 0$ implies $d\omega = 0$, and (ii) $d * \varphi = 0$ implies $d\psi_- = 0$. \square

A simple case is that in which the entire torsion τ_1 vanishes, which corresponds to

$$\begin{aligned} d\varphi &= \omega \wedge \rho, \\ d * \varphi &= -\psi_- \wedge \rho = -\psi_- \wedge \rho_2. \end{aligned}$$

This situation applies when M is the torus $T^6 = \mathbb{R}^6/\mathbb{Z}^6$ endowed with a constant $SU(3)$ -structure. The set (1) of such structures compatible with \hat{g} is isomorphic to the set of G_2 -structures compatible with (20) (assuming that orientations are also preserved). We deduce that \mathbb{M} is calibrated if and only if $\rho = 0$, and cocalibrated if and only if $\rho_2 = 0$.

Examples. 1. Let \mathbb{M} be a circle bundle with curvature 2-form

$$\rho = e^{12} + e^{34} + e^{56}$$

over T^6 . Then there are no calibrated G_2 structures on \mathbb{M} compatible with g . The set \mathcal{C}^* of cocalibrated structures corresponds to $SU(3)$ -structures relative to which ρ has type $(1, 1)$. By considering first the space \mathbb{CP}^3 (regarded as the set $SO(6)/U(3)$ of orthogonal almost complex structures^{18,19}), \mathcal{C}^* can be seen to be a disjoint union $\mathbb{RP}^1 \sqcup \mathbb{RP}^5$.

2. We can apply the above theory to the real projective space \mathbb{RP}^7 by representing it as $SO(5)/SU(2)$, which fibres over $SO(5)/U(2) \cong \mathbb{CP}^3$. It is well known that the latter has a homogeneous nearly-Kähler metric, so that there is a reduction to $SU(3)$ with $\tau_1 \in \mathcal{W}_1^\pm$. Moreover, the curvature 2-form ρ of the above Hopf fibration is an element of $[\Lambda_0^{1,1}]$ relative to the non-integrable complex structure.²⁰ Example 1 of Section 3 now implies that \mathbb{RP}^7 has a G_2 -structure with $\tau_2 \in \mathcal{X}_1 \oplus \mathcal{X}_3$ and another with $\tau_2 \in \mathcal{X}_3 \oplus \mathcal{X}_4$.

3. Let us return to the descriptions (1). The space of G_2 -invariant differential forms on \mathbb{RP}^7 is 1-dimensional, and it follows that \mathbb{RP}^7 has a nearly-parallel G_2 -structure, one with $\tau_2 \in \mathcal{X}_1$. More generally, we may regard (16) as a list of $SO(6)$ -invariant differential forms on \mathbb{RP}^7 subject to the relations $d\alpha = -\frac{1}{2}\omega$ and

$$d\psi_{\pm} = \alpha \wedge \psi_{\mp}, \quad (22)$$

that also give rise to G_2 -structures with $\tau_2 \in \mathcal{X}_1 \oplus \mathcal{X}_3$.

One can arrive at (22) by considering the canonical S^1 -bundle \mathbb{B} over a Kähler manifold M of real dimension 6 (see (4)). A local orthonormal basis $\{\psi^1, \psi^2\}$ of sections of the bundle with fibre $[\Lambda^{3,0}] \cong \mathbb{R}^2$ gives rise to coordinates a^1, a^2 and radial parameter $r = \sqrt{(a^1)^2 + (a^2)^2}$ on \mathbb{B} . The Kähler condition implies that $d\psi^1 = \sigma \wedge \psi^2$ for some 1-form σ on M , and we use

$$b^1 = da^1 - a^2\pi^*\sigma, \quad b^2 = da^2 + a^1\pi^*\sigma$$

to define global forms

$$\begin{aligned} r dr &= a^1 b^1 + a^2 b^2, & \alpha &= a^1 b^2 - a^2 b^1, \\ \psi_+ &= a^1 \psi^1 + a^2 \psi^2, & \psi_- &= a^1 \psi^2 - a^2 \psi^1. \end{aligned}$$

A straightforward calculation² shows that

$$r dr \wedge \psi_+ + \alpha \wedge \psi_- = r^2(b^1 \wedge \psi^1 + b^2 \wedge \psi^2) = r^2 d\psi_+,$$

so that restricting to \mathbb{B} ($r = 1$) we obtain (22). Similarly, $d\alpha = \pi^*\rho$, where $\rho = d\sigma$ can be identified with the Ricci form. This leads to the following result of Baum et al.²¹ which forms part of a more general theory of nearly-parallel G_2 structures.^{22,15}

5.2 Theorem If M is Kähler-Einstein with positive scalar curvature then \mathbb{B} has a G_2 -structure with $\tau_2 \in \mathcal{X}_1$.

We conclude this article by returning to the initial set-up of this section, together with the assumption that the holonomy of the metric (20) reduces to the subgroup G_2 determined by the 3-form (12). This allows us to set $X_i = 0$ in Theorem 5.1.

5.3 Corollary If the holonomy group of the metric (20) reduces to G_2 then the quotient $M = \mathbb{M}/S^1$ has an $SU(3)$ -structure for which $\tau_1 \in \mathcal{W}_2^+$.

Observe that the resulting condition on τ_1 involves a change of sign from that in Corollary 3.2. Indeed, from (21), we obtain $d\omega = 0$ and $d\psi_- = 0$. It is convenient to regard $SU(3)$ -structures with $(d\psi_-)^{2,2} = 0$ as ‘self-dual’, and those with $(d\psi_+)^{2,2} = 0$ as ‘anti-self-dual’. The latter type occurred naturally in Sections 3 and 4, and we focus attention on the $(2, 2)$ components so that the terminology is conformally invariant. In the present situation, we may therefore say that M has a self-dual symplectic structure with

$$d\psi_+ = \omega \wedge \rho,$$

where $\rho = \rho_1$ is a closed effective $(1, 1)$ -form.

Whilst the incomplete examples of Section 4 admit quotients of this type, a more realistic generalization of the condition $\tau_1 = 0$ is obtained by dropping the assumption that the S^1 orbits on \mathbb{M} have constant size. In this situation, there exists a function f on M for which $d(e^{2f}\omega) = 0$. It follows that, applying the conformal transformation (10), M once again has a self-dual symplectic structure, though this time $W_5 = -df$ is non-zero and

$$\rho = \rho_1 + \rho_2 = -e^{-3f}(W_2^+ + 2df \lrcorner \psi_-)$$

is closed. A study of this particular class of structures may be valuable in the construction of metrics with holonomy G_2 .

Acknowledgments

This paper was conceived for the conference ‘Differential Geometry Valencia 2001’, though the material was developed in the light of feedback from the Durham-LMS Symposium ‘Special Structures in Differential Geometry’ that took place a month later. The authors wish to thank the organizers of both conferences.

References

1. R.L. Bryant. Metrics with exceptional holonomy. *Annals of Math.*, 126:525–576, 1987.
2. S. Salamon. *Riemannian geometry and holonomy groups*. Pitman Research Notes in Mathematics 201. Longman, 1989.
3. N. Hitchin. The geometry of three-forms in six dimensions. *J. Differ. Geom.*, 55:547–576, 2000.
4. N. Hitchin. Stable forms and special metrics. In *Global Differential Geometry: The Mathematical Legacy of Alfred Gray*, volume 288 of *Contemp. Math.*, pages 70–89. American Math. Soc., 2001.

5. G. W. Gibbons, H. Lü, C. N. Pope, and K. S. Stelle. Supersymmetric domain walls from metrics of special holonomy. hep-th/0108191.
6. R. Bryant and S. Salamon. On the construction of some complete metrics with exceptional holonomy. *Duke Math. J.*, 58:829–850, 1989.
7. M. Atiyah and E. Witten. M-theory dynamics on a manifold of G_2 holonomy. hep-th/0107177.
8. L. Falcitelli, A. Farinola, and S. Salamon. Almost-Hermitian geometry. *Differ. Geom. Appl.*, 4:259–282, 1994.
9. A. Gray and L. Hervella. The sixteen classes of almost Hermitian manifolds. *Ann. Mat. Pura e Appl.*, 282:1–21, 1980.
10. M. Fernández and A. Gray. Riemannian manifolds with structure group G_2 . *Ann. Mat. Pura Appl.*, 32:19–45, 1982.
11. G. W. Gibbons, D. N. Page, and C. N. Pope. Einstein metrics on S^3 , R^3 , and R^4 bundles. *Commun. Math. Phys.*, 127:529–553, 1990.
12. D. D. Joyce. *Compact manifolds with special holonomy*. Oxford University Press, 2000.
13. M. Fernández. An example of a compact calibrated manifold associated with the exceptional Lie group G_2 . *J. Differ. Geom.*, 26:367–370, 1987.
14. F. M. Cabrera. On Riemannian manifolds with G_2 -structure. *Boll. Un. Mat. It.*, 10:99–112, 1996.
15. F. M. Cabrera, M. D. Monar, and A. F. Swann. Classification of G_2 -structures. *London Math. Soc.*, 53:407–416, 1996.
16. A. Gray. The structure of nearly Kähler manifolds. *Math. Ann.*, 223:233–248, 1976.
17. S. M. Salamon. Almost product structures. In *Global Differential Geometry: The Mathematical Legacy of Alfred Gray*, volume 288 of *Contemp. Math.*, pages 162–181. American Math. Soc., 2001.
18. E. Abbena, S. Garbiero, and S. Salamon. Almost Hermitian geometry of 6-dimensional nilmanifolds. *Ann. Sc. Norm. Sup.*, 30:147–170, 2001.
19. V. Apostolov, G. Grantcharov, and S. Ivanov. Orthogonal complex structures on certain Riemannian 6-manifolds. *Diff. Geom. Appl.*, 11:279–296, 1999.
20. R. Reyes-Carrion. A generalization of the notion of instanton. *Differ. Geom. Appl.*, 8:1–20, 1998.
21. H. Baum, Th. Friedrich, R. Grünewald, and I. Kath. *Twistor and Killing spinors on Riemannian manifolds*. Teubner-Verlag, Stuttgart-Leipzig, 1991.
22. Th. Friedrich, I. Kath, A. Moroianu, and U. Semmelmann. On nearly parallel G_2 structures. *J. Geom. Phys.*, 27:155–177, 1998.